

Claremont Colleges Scholarship @ Claremont

All HMC Faculty Publications and Research

HMC Faculty Scholarship

12-1-2001

A Local Inversion Principle of the Nash-Moser Type

Alfonso Castro
Harvey Mudd College

J. W. Neuberger
University of North Texas

Recommended Citation

Castro, Alfonso and Neuberger, J. W., "A Local Inversion Principle of the Nash-Moser Type" (2001). *All HMC Faculty Publications and Research*. Paper 479.
http://scholarship.claremont.edu/hmc_fac_pub/479

This Article is brought to you for free and open access by the HMC Faculty Scholarship at Scholarship @ Claremont. It has been accepted for inclusion in All HMC Faculty Publications and Research by an authorized administrator of Scholarship @ Claremont. For more information, please contact scholarship@cuc.claremont.edu.

A LOCAL INVERSION PRINCIPLE OF THE NASH–MOSER TYPE*

ALFONSO CASTRO[†] AND J. W. NEUBERGER[‡]

Abstract. We prove an inverse function theorem of the Nash–Moser type. The main difference between our method and that of [J. Moser, *Proc. Nat. Acad. Sci. USA*, 47 (1961), pp. 1824–1831] is that we use continuous steepest descent while Moser uses a combination of Newton-type iterations and approximate inverses. We bypass the *loss of derivatives problem* by working on finite dimensional subspaces of infinitely differentiable functions.

Key words. Nash–Moser methods, inverse function theorem, continuous steepest descent

AMS subject classifications. Primary, 34B15; Secondary, 35J65

PII. S0036141099357690

1. Introduction. Inverse function theorems are fundamental tools for the study of solutions to nonlinear equations. Proofs depend on iteration arguments. When a nonlinear equation comes from a partial differential equation, it often happens that the operators under consideration do not have enough *regularity* and the iteration process is defined only for a few steps. This is known as loss of derivatives. In order to overcome this difficulty, iteration techniques have been designed to allow the iteration to lead to a limit. These are known as generalized inverse function theorems. They generally depend on having a *scale* of intermediate spaces between the domain and the range of the nonlinear operator under consideration. Most notable of such results is the one proven by J. Moser in [5] and [6] (see also [2]). The reader is referred to [3] for a sharper version of the results of [5].

Here we prove an inverse function theorem (Theorem 2.1 below) using finite dimensional subspaces of infinitely differentiable functions, which makes the phenomenon of loss of derivatives immaterial. In addition we only assume the operators to have a first order derivative. This is in contrast with the results of [5] and [3] where Newton-like iteration techniques require extensive use of the properties of the *quadratic remainder* (see (10)). In [12] another inverse function theorem is proven without assumptions on the quadratic remainder. Our use of continuous steepest descent has roots in [1], [7], [8], and [9]. For applications of generalized inverse function theorems to elliptic systems the reader is referred to [4] and [11].

2. Main result. For the sake of simplicity we present our main result in the context of a particular class of Sobolev spaces. However, the general principle applies to many other *scales* of spaces sharing the general properties of the Sobolev spaces that are defined in the next paragraph. The reader is referred to [2] for various other scales of spaces.

Let each of m and n denote a positive integer. For each nonnegative integer ρ let C^ρ denote the set of ρ -differentiable functions $u : R^n \rightarrow R^m$ that are 2π -periodic in each of its n independent real variables. The norm in C^ρ is given by

$$(1) \quad |u|_\rho = \max\{|u(x)|; x \in R^n\} + \max\{|D^\alpha u(x)|; x \in R^n, |\alpha| = \rho\};$$

*Received by the editors June 23, 1999; accepted for publication (in revised form) June 7, 2001; published electronically December 28, 2001.

<http://www.siam.org/journals/sima/33-4/35769.html>

[†]Division of Mathematics and Statistics, University of Texas at San Antonio, San Antonio, TX 78249-0664 (acastro@utsa.edu).

[‡]Department of Mathematics, University of North Texas, Denton, TX 76203 (jwn@unt.edu).

here, $\alpha = (\alpha_1, \dots, \alpha_n)$ is an n -tuple of nonnegative integers and $|\alpha| = \alpha_1 + \dots + \alpha_n$. Similarly for $r \geq 0$ we define H^r as the Sobolev space of functions $u : R^n \rightarrow R^m$ of the form

$$(2) \quad u(x) = \sum_{j \in \mathbf{Z}^n} c_j e^{ij \cdot x}$$

such that

$$(3) \quad \|u\|_r^2 \equiv \sum_{j \in \mathbf{Z}^n} |c_j|^2 + \sum_{j \in \mathbf{Z}^n} |j|^{2r} |c_j|^2 < \infty,$$

where $|(j_1, \dots, j_n)|^2 = j_1^2 + \dots + j_n^2$. Actually H^r is also defined for $r < 0$ provided that the expression in (2) is understood in the sense of distributions and

$$(4) \quad \|u\|_r^2 \equiv |c_0|^2 + \sum_{j \in \mathbf{Z}^n - \{0\}} |j|^{2r} |c_j|^2 < \infty.$$

For $r \geq 0$ the inner product in H^r is given by

$$(5) \quad \left\langle \sum_{j \in \mathbf{Z}^n} c_j e^{ij \cdot x}, \sum_{j \in \mathbf{Z}^n} d_j e^{ij \cdot x} \right\rangle_r = \sum_{j \in \mathbf{Z}^n} (1 + |j|^{2r}) c_j \bar{d}_j.$$

Let $\epsilon > 0$ and $F : \{x \in C^1; |x|_1 < \epsilon\} \rightarrow C^0$ be a continuous function such that $F(0) = 0$. Typically F is a first order differential operator. We assume that for $u, v \in C^1$, with $|u|_1 < \epsilon$, the limit

$$(6) \quad \lim_{t \rightarrow 0} \frac{F(u + tv) - F(u)}{t} \equiv F'(u)v$$

exists and is a continuous function of v and that $F'(\cdot)v$ defines a continuous function from C^1 into C^0 , for each $v \in C^1$. We also assume that there exist $\lambda \in R$, positive constants k_1, k_2, k_3 , and $l > (n/2) + 1$ with

$$(7) \quad \langle F'(u)v, v \rangle_0 \geq k_1 \|v\|_0^2$$

for all $u, v \in C^1$;

$$(8) \quad \langle F'(u)v, v \rangle_l \geq k_2 \|v\|_\lambda^2 - k_3 \|v\|_0^2$$

for $u \in C^1$, $|u|_l \leq \epsilon$, $v \in H^l$; and for each positive integer ρ there exists M_ρ such that

$$(9) \quad \|F(u)\|_\rho \leq M_\rho \|u\|_{\rho+1} \quad \text{for } |u|_1 < \epsilon.$$

Without loss of generality we may assume that $k_1 < k_2$.

In [6], it is shown that the equation $F(u) = g$ has a solution when the operators $F'(u)$ have *approximate inverses*, and

$$(10) \quad \|F(u+v) - F(u) - F'(u)v\|_0 \leq M \|v\|_0^{2-\beta} \|v\|_l^\beta \quad \text{for } |u|_1 < \epsilon, |u+v|_1 < \epsilon,$$

where M is a constant independent of u, v and $0 \leq \beta < 1$. In addition, it is shown that (7) and (8) with $l = \lambda$ are sufficient for $F'(u)$ to have an approximate inverse.

Here we prove the following theorem.

THEOREM 2.1. *If (6)–(9) hold and $l > (n/2) + 1$, then there exist $\delta > 0$ such that, if $\|g\|_l < \delta$, then the equation*

$$(11) \quad F(y) = g$$

has a solution.

Proof. For each positive integer k , let X_k denote the linear subspace of H^l of functions of the form

$$(12) \quad u(x) = \sum_{\|j\|^2 \leq k^2} c_j e^{ij \cdot x}.$$

Let $P_k \equiv P$ denote the orthogonal projection of H^l onto X_k . An elementary Fourier series argument shows that P is also an orthogonal projection of H^0 onto X_k and

$$(13) \quad \langle Pu, v \rangle_0 = \langle u, Pv \rangle_0 \quad \text{for all } u, v \in H^l.$$

Now let $\Lambda \in (n/2 + 1, l)$. Since X_k is finite dimensional and a subset of C^1 , by (6) there exists a bounded differentiable function $\tilde{F} : X_k \rightarrow X_k$ such that $PF(u) = \tilde{F}(u)$ if $\|u\|_\Lambda < \epsilon/2$, $u \in X_k$. Hence the initial value problem

$$(14) \quad z'(t) = -\tilde{F}(z(t)) + P(g), \quad t \geq 0, \quad z(0) = 0,$$

has a solution defined on $[0, \infty)$.

Let us see that, for $\|g\|_l$ small enough, $|z(t)|_1 < \epsilon/2$ for all $t \geq 0$. In fact, let $w(t) = P(\tilde{F}(z(t)) - g)$. Thus

$$\begin{aligned} (\|w(t)\|_0^2)' &= 2\langle w(t), P\tilde{F}'(z(t))z'(t) \rangle_0 \\ (15) \quad &= -2\langle Pw(t), \tilde{F}'(z(t))(P(w(t))) \rangle_0 \\ &\leq -2k_1\|w(t)\|_0^2. \end{aligned}$$

In particular we see that the quantity $\|w(t)\|_0$ is a decreasing function of t . In addition, from (15) we have

$$(16) \quad \|w(t)\|_0 \leq \|w(0)\|_0 e^{-k_1 t} = \|g\|_0 e^{-k_1 t}.$$

Now we estimate the H^l norm of $w(t)$. In order to do so we observe that for each (k, λ) there exists a positive constant $C(k, \lambda)$ such that

$$(17) \quad \|x\|_\lambda^2 \geq C(k, \lambda)\|x\|_l^2 \quad \text{for all } x \in X_k.$$

We note that $C(k, \lambda) \rightarrow 0$ as $k \rightarrow \infty$ when $\lambda < l$; otherwise $C(k, \lambda)$ can be taken to be equal to 1. From now on we restrict ourselves to the case $\lambda < l$; the case $\lambda \geq l$ is simpler. Thus we may assume that

$$(18) \quad C(k, \lambda) \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

From (8), (13), (16), (17), and (18) we infer that for k sufficiently large

$$\begin{aligned} (\|w(t)\|_l^2)' &= 2\langle w(t), P\tilde{F}'(z(t))z'(t) \rangle_l \\ (19) \quad &= -2\langle Pw(t), \tilde{F}'(z(t))(P(w(t))) \rangle_l \\ &\leq -2(k_2\|w(t)\|_\lambda^2 - k_3\|w(t)\|_0^2) \\ &\leq -2(k_2C(k, \lambda)\|w(t)\|_l^2 - k_3\|g\|_0^2 e^{-2k_1 t}). \end{aligned}$$

Thus for k sufficiently large

$$\begin{aligned}
 \|w(t)\|_l^2 &\leq (\|g\|_l^2 + k_3\|g\|_0^2/(k_1 - k_2C(k, \lambda)))e^{-2k_2C(k, \lambda)t} \\
 (20) \qquad &\leq (\|g\|_l^2 + 2k_3\|g\|_0^2/k_1)e^{-2k_2C(k, \lambda)t} \\
 &\equiv M(\|g\|_l)e^{-2k_2C(k, \lambda)t},
 \end{aligned}$$

where $M(\|g\|_l) \rightarrow 0$ as $\|g\|_l \rightarrow 0$. By interpolation properties of Sobolev space (see section I.2 in [6]) one has

$$(21) \quad \|w(t)\|_\Lambda \leq \|w(t)\|_0^{(1-\Lambda/l)} \|w(t)\|_l^{\Lambda/l} \leq M(\|g\|_l)^{\Lambda/(2l)} \|g\|_0^{(1-\Lambda/l)} e^{-k_1(1-\Lambda/l)t}.$$

Integrating (14) by (20) and (21), we see that there exist $\delta > 0$ such that if $\|g\|_l \leq \delta$, then

$$(22) \qquad \|z(t)\|_\Lambda < \epsilon/3 \quad \text{for all } t \geq 0.$$

Now letting $x_k \in X_k$ be an element in the w -limit set of the orbit defined by z , we see that $\|x_k\|_\Lambda \leq \epsilon/3$ and $\tilde{F}(x_k) = PF(x_k) = P(g)$. Therefore by the Sobolev imbedding theorem we may assume that it converges to some element $x \in H^{\lambda_1}$, with $\lambda_1 \in ((N/2) + 1, l)$. By (9), $F(x_k)$ is bounded in H^{λ_1-1} . Using again that bounded sequences in H^s have converging subsequences in H^t if $s > t$, we may further assume that $\{F(x_k)\}$ converges in H^{λ_2} with $\lambda_2 \in (n/2, \lambda_1 - 1)$. Recalling that, by Poincaré's inequality,

$$(23) \qquad \|z\|_{\lambda_2-1} \leq C_k \|z\|_{\lambda_2} \quad \text{for all } z \in X_k^\perp,$$

with C_k converging to zero as $k \rightarrow \infty$, we conclude that $\|(I - P)F(x_k)\|_{\lambda_2-1} \rightarrow 0$ as $k \rightarrow \infty$. This and the fact that $\{F(x_k)\}$ converges to $F(x)$ imply that $\{P(F(x_k))\}$ converges to $F(x)$ in H^{λ_2} . Hence in H^{λ_2} we have

$$(24) \qquad g = \lim P_k(g) = \lim P_k(F(x_k)) = F(x),$$

and this proves the theorem. \square

REFERENCES

- [1] A. CASTRO AND J. W. NEUBERGER, *An inverse function theorem*, Contemp. Math., 221 (1999), pp. 127–132.
- [2] R. HAMILTON, *The inverse function theorem of Nash and Moser*, Bull. Amer. Math. Soc. (N.S.), 7 (1982), pp. 65–222.
- [3] L. HÖRMANDER, *On the Nash-Moser implicit function theorem*, Ann. Acad. Sci. Fenn. Ser. A. I. Math., 10 (1985), pp. 255–259.
- [4] P. KORMAN, *On existence of solutions to two classes of nonlinear problems*, Comm. Partial Differential Equations, 14 (1989), pp. 519–539.
- [5] J. MOSER, *A new technique for the construction of solutions of nonlinear differential equations*, Proc. Nat. Acad. Sci., USA, 47 (1961), pp. 1824–1831.
- [6] J. MOSER, *A rapidly convergent iteration method and non-linear differential equations. I.*, Ann. Scuola Norm. Sup. Pisa (3), 20 (1966), pp. 265–315.
- [7] J. W. NEUBERGER, *Steepest descent and differential equations*, J. Math. Soc. Japan, 37 (1985), pp. 187–195.
- [8] J. W. NEUBERGER, *Constructive variational methods for differential equations*, Nonlinear Anal., 13 (1989), pp. 413–428.

- [9] J. W. NEUBERGER, *Sobolev Gradients and Differential Equations*, Lecture Notes in Math. 1670, Springer-Verlag, Berlin, 1997.
- [10] M. POPPENBERG, *An inverse function theorem for Fréchet spaces satisfying smoothing property and (DN)*, Math. Nachr., 206 (1999), pp. 123–145.
- [11] P. RABINOWITZ, *A rapid convergence method for a singular perturbation problem*, Ann. Inst. H. Poincaré Anal. Non Linéaire, 1 (1984), pp. 1–17.
- [12] W. O. RAY, *A rapidly convergent iteration method and Gâteaux differentiable operators*, J. Math. Appl., 103 (1984), pp. 162–171.